ISSN: 2792-8268

Volume: 38, Jan-2025

http://sjii.indexedresearch.org

# Potential Diagnostic Markers of Mycobacterium Tuberculosis Infection: A Bibliometric Study (2019-2023)

## Sabarina Elfrida Manik

Doctoral Program in Biomedicine, Postgraduate School, Universitas Yarsi

#### Himmi Marsianti

Doctoral Program in Biomedicine, Postgraduate School, Universitas Yarsi

### **Bobby Singh**

Doctoral Program in Biomedicine, Postgraduate School, Universitas Yarsi

### Wening Sari

Doctoral Program in Biomedicine, Postgraduate School, Universitas Yarsi

#### Inas Hasna Azizah

Faculty of Health and Technology, Binawan University

Abstract: This bibliometric analysis was conducted using the Scopus database. Many analysis, including those for the most prosperous nations, journals, authors, institutions, and articles, were performed on the papers that were retrieved. The existing condition, new areas of interest, and potential for chemokine and tuberculosis research were ascertained using co-occurrence mapping of phrases and keywords. VOSviewer and Bibliometrix were utilized to visualize the collaborative network mapping and evaluate the data. The results: The most common keywords were "chemokine" and "tubercukosis". The Scopus search engine identifed 802 Tuberculosis-related documents published by Global Countries between 2019 to 2023. According to our findings, United States was the most productive in Global Countries. In the early stages of the study, researchers investigated immune regulatory and signalling pathway in tuberculosis-related study. In recent years, the development of antivirus agents has emerged as a prominent topic. Conclusions: The study performed a bibliometric analysis of over 5 years of chemokine and tuberculosis research, identifying the nations, institution journal and publications active in this subject. The findings provide a comprehensive overview of chemokine in tuberculosis research.

**Keywords:** Tuberculosis, Bibliometric analysis, chemokine

#### Introduction

Tuberculosis (TB) is one of the ten causes of death and the main cause of infectious agents. About 1.3 million fatalities among HIV-negative individuals and approximately 300,000 deaths among HIV-positive individuals are attributable to tuberculosis. The 2023 World Health Organization (WHO) report gives a thorough and current evaluation of tuberculosis (TB) and the advancements made in the diagnosis, treatment, and prevention of this illness at the national, international, and local levels. This is

ISSN: 2792-8268

Volume: 38, Jan-2025

http://sjii.indexedresearch.org

done within the framework of TB targets, plans, and commitments (1). The report's 2023 edition is based on information gathered by WHO from national health ministries as part of its yearly data collection. More than 99% of the world's population lives in 192 nations and territories, and statistics on tuberculosis cases is reported for 2023 (2).

The bacterium Mycobacterium tuberculosis is the main cause of the infectious disease tuberculosis (MTB) which can attack various organs in the body except nails, hair or feathers and most often attacks the lungs because the initial entry of the bacteria is inhalation. This disease, if not treated properly, can cause various complications and even death (3). Despite scientific advances in The detection and management of tuberculosis continues to be the biggest hazard to public health, particularly in developing nations. It is estimated that latent tuberculosis infection, in which an intracellular bacterium infects an individual without causing an active disease but still puts them at risk of becoming active again, affects one-third of the world's population. Approximately 10% of people with primary MTB infections have the potential to become active. Most of the other MTB are found in the lungs in structures known as granulomas, where they are latent (4).

Currently, latent tuberculosis infection (ITBL) can be detected using the interferon-gamma release assay (IGRA). This examination can only determine whether an individual has been exposed to TB germs called other tuberculosis, but does not help differentiate TBL infection from active TB disease. The principle of IGRA is to detect Interferon-gamma (IFN-γ) which is excreted by T cells as a response to stimulation of specific Mtb antigens. A cohort study with a large sample size in a country with a high TB infection rate comparing the effectiveness of IGRA and the tuberculin test showed that no significant differences were found. To diagnose latent tuberculosis infection, tuberculin test, and QFT-GIT. This is supported by other cohort studies with the same results, there is no significant difference in predicting adult TB infection in South Africa. IGRA sensitivity detects 75% positive TB and tuberculin test 76.9%, the effectiveness of both is less than 50%: IGRA 49.3% and tuberculin test 45.0%. This is a consideration in assessing the effectiveness of the IGRA test compared to the relatively cheaper tuberculin test (4).

Recent research as an alternative for TB diagnosis has identified a chemokine, namely interferongamma inducible protein 10 (IP10) (5). IP10 is a chemokine, namely interferon, which is excreted by cells that present antigens in response to IFN-y by interacting with the CXCR3 receptor to control T cell migration to inflammatory areas. IP10 was found to increase in day 1 plasma that was not stimulated by adults and children suffering from TB (5). In HIV-infected patients the IP10 response is known to be greater and stronger than IFN-y (6). IP10 can also be detected in patients with active TB and its levels decrease after receiving therapy (7). Patients with systemic symptoms, such as fever and anorexia, had higher levels of IP10 (8). The results of research on IP10 in patients infected with TB, both adults and children, were higher than controls, thus showing potential value for diagnosing latent TB (9). Sodium electrolyte examination is one of the examinations that plays an important role in TB patients which indicates an electrolyte imbalance between those receiving TB treatment (5). Examination of the neutrophil lymphocyte ratio (NLR) percentage value can play a role as an immune response in TB sufferers, because the neutrophil response will increase after 8-15 days of being infected with Mtb and then last until the end of the infection. These results provide an understanding of how important neutrophils are in the early phase of infection. The adaptive immune system against Mtb mainly relies on CD4+ cells, the role of CD4 which produces the cytokine interferon gamma (IFN-γ) in the incidence of primary resistance to Tuberculosis (TBC) has been successfully identified (10). Thoracic x-ray examination is very important for diagnosing TB patients. The x-ray results can show infiltrative spots

ISSN: 2792-8268

Volume: 38, Jan-2025

http://sjii.indexedresearch.org

or nodules, especially in the upper lobes. Apart from that, thoracic x-rays can also show the formation of cavities, calcified nodules such as tubrtclouma and many small nodular lesions indicating Miler Tuberculosis (TBC). Radiological examination is very important for the diagnosis of Tuberculosis (TB). The presence of suggestive clinical symptoms of Tuberculosis (TB) is almost always an abnormality found on X-rays. If clinically supportive but X-ray radiological examination >10 weeks post-infection is negative, then this is a strong sign that what is being suffered is not tuberculosis (11). Diagnosis currently still experiences many obstacles, especially collecting sputum specimens which do not comply with standards, especially in collecting sputum from elderly patients and children so that the results obtained are not optimal.

Analyzing the general trend of research activity and examining the relationships between pertinent research institutions can be done effectively with bibliometric studies. The bibliometric studies can assess the volume and trends of scientific output in important biomedical fields across nations and years. They are especially helpful for new fields whose effects on the broader field of biomedical research are still being completely assessed (32). We present this study on the scientific productions in this subject among countries over the last five years to better understand chemokine's boundaries and hot regions in tuberculosis research (2019-2023).

#### **METHOD**

This research design is identification of research journal within 5 years, namely from 2019 to 2023. The research journal are those contain tuberculosis in Global Countries.

### Sources of information and techniques for searching

For this study, all of the data were obtained from the Web of Scopus. To prevent variations, literature retrieval was done all in one day on January 1st, 2024, taking into account the speed at which databases are updated. This study was published over a four-decade span. The following were the search phrases used: TS stands for [interleukine OR tuberculosis]. Only original articles and reviews published in English were included among the several publication types.

### Data collecting and cleaning

Comprehensive publishing parameters, such as the quantity of papers and citations, the H-index, the publication year, the nation or area, the affiliations, the authors, the journal, the references, and the keywords, were collected from the database for the current study. Spelling mistakes and duplicate writers were then eliminated. We believe that the majority of the original data are reliable, even though errors in the analysis may not be entirely prevented due to variations in the formats of referenced journals, different versions of cited references, and the same abbreviated name for several authors. Prior to data analysis using VOSviewer v.1.6.10.0 (Science and Technology Research Center, Leiden University, Leiden, the Netherlands), spelling mistakes were fixed, superfluous words were eliminated, and some repetitive terms were combined into one word. Lastly, for bibliometric analysis, cleaned data were imported into VOSviewer and the Bibliometrix software.

### Bibliometric analysis

The two primary criteria for assessing the caliber of research are productivity, which is generally assessed by the number of publications, and impact, which is measured by the number of citations, with the exception of self-citation. The Hindex has gained popularity recently as a tool for assessing academic contributions made by scholars and forecasting their future scientific successes. The H-index creates a threshold that connects the Np and Nc, combining productivity and effect. Put another way, a researcher will have an H-index if they have published H publications and each one has been referenced at least H

ISSN: 2792-8268

Volume: 38, Jan-2025

http://sjii.indexedresearch.org

times. Although the H-index was first designed to evaluate a student's academic performance, it can also be used to characterize the publication output of a nation or region, organization, or journal.

#### **RESULTS**

### **Overview of Publications on Tuberculosis**

Our study provided new prespective on the scientific contribution of global countries to tuberculosis related research from 2019 to 2023. the Scopus algorithm identified 802 documents published by Global countries in the tuberculosis related field. Our study yielded the following results: 561 (70.0%) articles, 200 (24.9%) reviews, 14 (1.7%) editorials, 9 (1.1%) letter, 6 (0.7%) book chapter, and 12 (1.5%) others. Of the aforementioned publications, the majority speak English (n=790; 22.3%).

Fig. 1 Trends of publication articles on tuberculosis and chemokine from 2019 to 2023 by Global Countries.

No.	Country	Number of Article	Percentage (%)
1.	United States	238	29,7%
2.	China	162	20,2%
3.	India	81	10,1%
4.	United Kingdom	78	9,7%
5.	South Africa	71	8,9%
6.	Germany	60	7,5%
7.	Netherlands	48	6,0%
8.	France	42	5,2%
9.	Italy	35	4,4%
10.	Australia	30	3,7%

The tuberculosis publishing trends in Global countries exhibited a gradual increase from 2019 to 2021. Tuberculosis and chemokine publishing trends globally exhibited a gradual increase from 1981 to 2023 (Fig. 1). In addition, the peak was 2021 published 220 following 2020 published 172 articles. The visualization of collaborating nations was shown in Figure 2. The relative size of the frames represents the number of collaborations. For example, the frame size of the United States is the largest among countries, in addition United Kingdom is larger than Germany, indicating that the United States (n=24) builds collaborative studies with most countries compared to other countries, following United Kingdom (n=23) builds more countries than Ghana (n=21).

The productivity among Global countries in research documents related to tuberculosis and chemokine was also ranked, with United States having the highest percentage. The top ten journals publishing articles related to chemokine in tuberculosis in Global countries were also listed (Table 2).

Table 2. The most cited tuberculosis-interleukin related papers published by Global countries

No	Author	Title	Journal	Year	Citation
----	--------	-------	---------	------	----------

ISSN: 2792-8268

Volume: 38, Jan-2025

http://sjii.indexedresearch.org

1.	Zhao, H., Wu, L., Yan, G.,Wu, Y., Li, Y.	progression: signaling	Signal Transduction and Targeted Theraphy, 6(1): 263	2021	804 (22)
2.	Soehnlein, O., <u>Libby, P.</u>	Targeting inflammation in atherosclerosis from experimental insights to the clinic	Nature Reviews Drug Discovery, 20(8), pp. 589-610	2021	485 (23)
3.	Batool, M., Ahmad, B., Choi, S.	A structure-based drug discovery paradigm	International Journal of Molecular Sciences, 20(11), 2783	2019	342 (24)
4.	Shah, V.K., Firmal, P., Alam, A., Ganguly, D., Chattopadhy ay, S.	Overview of Immune Response During SARS-CoV- 2 Infection: Lessons From the Past	Overview of Immune Response During SARS-CoV-2 Infection: Lessons From the Past	2020	326 (25)
5.	Fischinger, S., Boudreau, C.M., Butler, A.L., Streeck, H., Alter, G.	Sex differences in vaccine- induced humoral immunity	Seminars in Immunopathology 41(2), oo. 239-249	2019	247 (26)
6.	Mahesh, G., Biswas,R	MicroRNA-155: A Master Regulator of Inflammation	Journal of Interferon and Cytokine Research, 39(6), pp. 321-330	2019	200 (27)
7.	MacLean, E., <u>Broger</u> , T., <u>Yerlikaya</u> , S., <u>Pai</u> , M., <u>Denkinger</u> , C.M.	A systematic review of biomarkers to detect active tuberculosis	Nature Microbiology, 4(5), pp. 748-758	2019	137 (28)
8.	Ardain, A., Domingo- Gonzalez, R., Das, S.,Leslie, A., Khader, S.A.	Group 3 innate lymphoid cells mediate early protective immunity against tuberculosis	Nature, 570 (7762): 528-532	2019	135 (29)
9.	Pidwill, G.R., Gibson, J.F., Cole, J., Renshaw, S.A., Foster, S.J.	The Role of Macrophages in Staphylococcus aureus Infection	Frontiers in Immunology, 11, 620339	2021	134 (30)
10.	Alharbi, K.S., Fuloria, N.K., Fuloria, S.,Dua, K., Gupta, G.	Nuclear factor-kappa B and its role in inflammatory lung disease	Chemico-Biological Interactions, 345, 109568	2021	129 (31)

Volume: 38, Jan-2025

http://sjii.indexedresearch.org

The most documents were published at Frontiers In Immunology (n= 147; 18.3%), followed by Tuberculosis (n= 20; 2.5%), and Frontiers In Microbiology (n= 20; 2.5%). Also, the characteristics of papers published by Global countries with the highest number of citations in the last 5 years were listed in Table 3.

Table 3. Jurnal publishing articles related tuberculosis produced by Global countries


No.	Country	Number of Article	Percentage (%)
1.	Frontiers In Immunology	147	18,3%
2.	Tuberculosis	20	2,5%
3.	Frontiers In Microbiology	20	2,5%
4.	Scientific Reports	19	2,4%
5.	International Journal Of Molecular Sciences	18	2,2%
6.	Frontiers In Cellular And Infection Microbiology	17	2,1%
7.	Plos Pathogens	15	1,9%
8.	Cells	15	1,9%
9.	Vaccines	10	1,2%
10.	Pathogens	9	1,1%

An author's term network visualization from Global countries in 5 years period of tuberculosis related research was illustrated in Figure 3a. A total of 802 distinct types of publications were found in the retrieved papers, 25 of which appeared more than 100 times. The following is an evaluation of the primary three central clusters: Terms pertaining to "macrophage," "nonhuman," and "immune response" are found in cluster 1 (shown in red); cluster 2 (shown in green) contains "tuberculosis", "human", "male", "female"; and cluster 3 (in blue) related to "tumor necrosis factor", "cytokine", "interleukine 10", "gamma interferon". VOSviewer software also designed colors to chemokine tuberculosis related terms based on publication years (Figure 3b). Terms that first occur in the early years are shown by the color purple, whereas terms that have emerged recently are indicated by the color yellow. The development of chemokine research in tuberculosis demonstrates a dynamic shift over time, with an emphasis on "immunology", "metabolism" in 2020; "controlled study", "humans" interferon inducible pro" in 2021; "tuberculosis", "cd4+ t lymphocyte", "interleukin 6" in 2021. The institution from global

ISSN: 2792-8268

Volume: 38, Jan-2025

http://sjii.indexedresearch.org



nations with the highest rates of production in the chemokine tuberculosis from 2019-2023 were listed, and highlighting the prominent contributions of South Africa, US and UK.

### **DISCUSSION**

Our research offered a fresh viewpoint on the scientific contribution of Global countries to chemokine in tuberculosis related research. In general, US contribute 29.7% of the worldwide chemokine production in areas connected to TB. The quantity of articles released by international nations skyrocketed in 2020, rising from 128 in 2019 to 220 papers. Following 2021, chemokine tuberculosis-related sectors generated by international nations were noted, suggesting that tuberculosis-related issues are becoming more and more important as concerns for human health.

In 2019, Unites States was contributes the most article with 40 documents. Article in 2019 most discussed the biomarker and signaling pathways of tuberculosis. Meanwhile, Hawerkamp et al reportef IL-26 activated macrophage can killing of M. tuberculosis (12). Also, in 2020 began to develop research on the BCG vaccine (13) and immunocompetent cells as therapeutic targets in TB (14).

It might be challenging to determine the weight or effect of publications directly in bibliometric study. Nonetheless, other research contended that examining the connection between article relevancy and journal significance could yield significant insights (15). Furthermore, as Table 3 illustrates, the top ten most referenced articles may offer insights into contemporary patterns, the evolving field of study subjects, and significant avenues for further investigation.

After 2021, Research focuses more on the development of the BCG vaccine through molecular targets in TB, and analysis of cytokine responses as TB vaccine biomarkers (16)(17)(18). Effective antimicroba In order to lower mortality and counteract the increasing prevalence of Mtb infection, interventions are needed. Furthermore, it has been suggested and researched as an alternative to treat MTB infection by repurposing already existing medications from natural sources. Furthermore, insilico techniques have been used to identify potential compounds that combat MTB (18)(19)(20).

The global study have been reported that drugs to against MTB cannot effective since the existence of Multidrug resistant TB (MDR TB). Improper use of antibiotics causes TB bacteria to become resistant, making treatment less effective. In order to treat the infection, natural antimicrobial ingredients are needed (21).

http://sjii.indexedresearch.org

This study evaluates the trajectory and present status of research on tuberculosis published by international countries using bibliometric analysis. Nonetheless, certain limitations were noted that are also present in other bibliometric analyses. Because our research focused on articles indexed in Scopus, it's possible that our analysis missed any papers that contained the term "tuberculosis." Notwithstanding its limitations, this analysis is the first to offer a succinct summary of the chemokine in research related to tuberculosis that has been published by international nations.





Figure 1. Mapping of international cooperation amongst nations. Larger frame countries indicate comparatively more collaborative efforts.

Figure 2. A network visualization of Co-occurance netw ork of terms extracted from articles published by Global country in TB related study in 4 decades. The minimum number of occurances was set to 100 times.



Volume: 38, Jan-2025

http://sjii.indexedresearch.org




Figure 3. a network representation of the co-occurence network of phrases taken from papers that Global Country produced in a four-decade study related to tuberculosis. One hundred occurrences was the minimum quantity that was required (a); Visualization overlay of a network representation of the co-occurence network of phrases taken from papers that Global Country produced in a four-decade study related to tuberculosis. Blue denotes those that surfaced prior to those designated in yellow (b).

#### **CONCLUSION**

The objectives of the current study was to present an overview of the trend and new areas especially chemokine that has play role in the field of Tuberculosis disease. With 802 articles published in journals over the last five years, research production on Mycobaterium tuberculosis-related themes has improved in global countries. Prioritizing research on biomarkers and signaling pathways was done in the past. Nonetheless, study themes have evolved in recent years to examine vaccine target theraphy MTB. The largest number of publications pertaining to chemokine TB is attributed to the United States. The results of this study can be used as a guide for continuing research in developing nations related to tuberculosis.

#### **References:**

- 1. World Health Organization (2014) Global Tuberculosis Report 2014.
- 2. World Health Organization (2023) Global Tuberculosis Report 2023.
- 3. World Health Organization (2016) Global Tuberculosis Programme: Guidelines on the Management of latent Tuberculosis Infection
- 4. Zuniga J, Gracia DT and Mendoza TS (2012) "Cellular and Humoral Mechanisms Involved in the Control of Tuberculosis," Clinical and Developmental Immunology, pp. 1–18.
- 5. Petrone, L. et al. (2021) "Coinfection of Tuberculosis and COVID-19 Limits the Ability to in Vitro Respond to SARS-CoV-2.," Internation Journal of Infectious Diseases, 113, pp. 82–87.
- 6. Petrone, L. et al. (2018) "Evaluation of IP-10 in Quantiferon-Plus as Biomarker for the Diagnosis of Latent Tuberculosis Infection," Tuberculosis, 111, pp. 147–153.
- 7. A Cannas, L. Calvo and T Chiacchio (2010) "IP-10 Detection in Urine is Associated With Lung Diseases," BMC Infectious Diseases, 10.

ISSN: 2792-8268

Volume: 38, Jan-2025

http://sjii.indexedresearch.org

- 8. Andersen P et al. (2000) "Spesific Immunebased Diagnosis of Tuberculosis," Lancet, 356, pp. 104–1099.
- 9. Mori T et al. (2004) "Spesific Detection Of Tuberculosis Infection: An Interferongamma Based Assay Using New Antigens," Am J Respir Crit Care Med, 170(1), pp. 59–64.
- 10. Muttaqin, A. (2012) Buku Ajar Asuhan Keperawatan Klien dengan Gangguan Sistem Pernapasan. Jakarta: Salemba Medika.
- 11. Crofton (2002) Tuberculosis Klinik. 2nd ed. Jakarta: Widya Medika.
- 12. Hawerkamp, H.C., van Geelen, L., Korte, J. et al. Interleukin-26 activates macrophages and facilitates killing of Mycobacterium tuberculosis. Sci Rep 10, 17178 (2020). https://doi.org/10.1038/s41598-020-73989-y
- 13. Subbian S, Singh P, Kolloli A, Nemes E, Scriba T, Hanekom WA, Kaplan G. BCG Vaccination of Infants Confers Mycobacterium tuberculosis Strain-Specific Immune Responses by Leukocytes. ACS Infect Dis. 2020 Dec 11;6(12):3141-3146. doi: 10.1021/acsinfecdis.0c00696. Epub 2020 Nov 23. PMID: 33226778.
- Dallenga, T.K., Schaible, U.E. (2021). Neutrophil-Mediated Mechanisms as Targets for Host-Directed Therapies Against Tuberculosis. In: Karakousis, P.C., Hafner, R., Gennaro, M.L. (eds) Advances in Host-Directed Therapies Against Tuberculosis . Springer, Cham. https://doi.org/10.1007/978-3-030-56905-1\_13
- 15. Zhang S, Fan H, Zhang Y. 2021. "The 100 top-cited studies on dyslexia research: a Bibliometric analysis." Front Psychiatry. 12:714627. Frontiers Media SA.
- Bar-Oz M, Martini MC, Alonso MN, Meir M, Lore NI, Miotto P, Riva C, Angala SK, Xiao J, Masiello CS, Misiakou MA, Sun H, Moy JK, Jackson M, Johansen HK, Cirillo DM, Shell SS, Barkan D. The small non-coding RNA B11 regulates multiple facets of Mycobacterium abscessus virulence. PLoS Pathog. 2023 Aug 21;19(8):e1011575. doi: 10.1371/journal.ppat.1011575. PMID: 37603560; PMCID: PMC10470900.
- 17. Seth P, Dubey S. IL-22 as a target for therapeutic intervention: Current knowledge on its role in various diseases. Cytokine. 2023 Sep;169:156293. doi: 10.1016/j.cyto.2023.156293. Epub 2023 Jul 11. PMID: 37441942.
- 18. Zhuang L, Ye Z, Li L, Yang L, Gong W. Next-Generation TB Vaccine: Progress, Challenges, and Prospects. Vaccine 2023; 11(8): 1304; https://doi.org/10.3390/vaccines11081304
- 19. Khokhar M, Purohit P, Gadwal A, Tomo S, Bajpai NK, Shukla R
- 20. The Differentially Expressed Genes Responsible for the Development of T Helper 9 Cells From T Helper 2 Cells in Various Disease States: Immuno-Interactomics Study
- 21. JMIR Bioinform Biotech 2023;4:e42421
- 22. doi: 10.2196/42421
- 23. Irhan LM, Adikusuma W, Perwitasari DA. Genomic variants-driven drug repurposing for tubercolusosis by utilizing the established bioinformatics-based approach. Biochemistry and Biophysics Reports, 2022; 32: 101334. DOI: https://doi.org/10.1016/j.bbrep.2022.101334
- 24. Maiolini M, Gause S, Taylor J, Steakin T, Shipp G, Lamichhane P, Deshmukh B, Shinde V, Bishayee A, Deshmukh RR. The War against Tuberculosis: A Review of Natural Compounds and Their Derivatives. Molecules. 2020 Jun 30;25(13):3011. doi: 10.3390/molecules25133011. PMID: 32630150; PMCID: PMC7412169.

ISSN: 2792-8268

Volume: 38, Jan-2025

http://sjii.indexedresearch.org

- 25. Zhao H., Wu, L., Yan, G., ...Wu, Y., Li, Y. Inflammation and tumor progression: signaling pathways and targeted intervention. Signal Transduction and Targeted Theraphy, 2021;6(1): 263
- 26. Soehnlein, O., Libby, P. Targeting inflammation in atherosclerosis from experimental insights to the clinic. Nature Reviews Drug Discovery, 2021;20(8), pp. 589-610
- 27. Batool, M., Ahmad, B., Choi, S. A structure-based drug discovery paradigm. International Journal of Molecular Sciences, 2019; 20(11), 2783
- 28. Shah, V.K., Firmal, P., Alam, A., Ganguly, D., Chattopadhyay, S Overview of Immune Response During SARS-CoV-2 Infection: Lessons From the Past. Overview of Immune Response During SARS-CoV-2 Infection: Lessons From the Past 2020
- 29. Fischinger, S., Boudreau, C.M., Butler, A.L., Streeck, H., Alter, G. Sex differences in vaccine-induced humoral immunity. Seminars in Immunopathology 41(2), oo. 2019:239-249
- 30. Mahesh, G., Biswas, R. MicroRNA-155: A Master Regulator of Inflammation. Journal of Interferon and Cytokine Research, 2019;39(6), pp. 321-330
- 31. MacLean, E., Broger, T., Yerlikaya, S., ...Pai, M., Denkinger, C.M. A systematic review of biomarkers to detect active tuberculosis. Nature Microbiology, 2019;4(5), pp. 748-758
- 32. Ardain, A., Domingo-Gonzalez, R., Das, S., ...Leslie, A., Khader, S.A. Group 3 innate lymphoid cells mediate early protective immunity against tuberculosis. Nature, 2019;570 (7762): 528-532
- 33. Pidwill, G.R., Gibson, J.F., Cole, J., Renshaw, S.A., Foster, S.J. The Role of Macrophages in Staphylococcus aureus Infection. Frontiers in Immunology, 2021;11, 620339
- 34.Alharbi, K.S., Fuloria, N.K., Fuloria, S., ...Dua, K., Gupta, G. Nuclear factor-kappa B and its role in inflammatory lung disease, 2021 Chemico-Biological Interactions, 345, 109568.